

Sub-eV upper limits on neutrino masses from cosmology

Øystein Elgarøy*

Institute of theoretical astrophysics, University of Oslo, P.B. 1029 Blindern, 0315 Oslo, Norway

Ofer Lahav†

*Department of Physics and Astronomy, University College London,
Gower Street, London WC1E 6BT, United Kingdom*

(Dated: February 2, 2008)

Upper limits on neutrino masses from cosmology have been reported recently to reach the impressive sub-eV level, which is competitive with future terrestrial neutrino experiments. In this brief review of the latest limits from cosmology we point out some of the caveats that should be borne in mind when interpreting the significance of these limits.

I. INTRODUCTION

The latest results from the WMAP satellite [1] confirm the success of the Λ CDM model, where $\sim 75\%$ of the mass-energy density is in the form of dark energy, with matter, most of it in the form of cold dark matter (CDM) making up the remaining 25%. Neutrinos with masses on the eV scale or below will be a hot component of the dark matter and will free-stream out of overdensities and thus wipe out small-scale structures. This fact makes it possible to use observations of the clustering of matter in the universe to put upper bounds on the neutrino masses. An excellent review of the subject is that of Lesgourgues and Pastor [3]. With the improved quality of cosmological data sets seen in recent years, the upper limits have improved, and some quite impressive claims have been made in the recent literature. We will in the following summarize the latest upper bounds and point out some of the potential systematic uncertainties that need to be clarified in the future.

II. CURRENT CONSTRAINTS ON NEUTRINO MASSES

One of the assumptions underlying cosmological neutrino mass bounds is that they have no non-standard interactions and that they decouple from the thermal background at temperatures of order 1 MeV. In that case, the relation between the sum of the neutrino masses M_ν and their contribution to the energy density of the universe is given by $\Omega_\nu h^2 = M_\nu/93.14$ eV [2], where h is the dimensionless Hubble parameter defined by writing the Hubble constant as $H_0 = 100h$ km s $^{-1}$ Mpc $^{-1}$. The limits in table I show an impressive progression, and with limits reaching deep into the sub-eV region it is prudent to point out that there are systematic uncertainties associated with the limits. We will in the following discuss

briefly two different types of uncertainties: cosmological uncertainties (models and priors) and astrophysical uncertainties (e.g. galaxy-dark matter bias).

Data	Authors	M_ν -bound
2dFGRS (P01)	Elgarøy et al. 02[4]	1.8 eV
CMB+2dFGRS(C05)	Sanchez et al. 05[5]	1.2 eV
CMB+LSS+SNIa+BAO	Goobar et al. 06[6]	0.62 eV
WMAP (3 year) alone	Fukugita et al. 06[7]	2.0 eV
CMB+LSS+SN +	Spergel et al. 06[1]	0.68 eV
CMB + LSS+SNIa+BAO+Ly α	Seljak et al. 06[8]	0.17 eV

TABLE I: Some recent cosmological neutrino mass bounds (95 % CL).

III. COSMOLOGICAL UNCERTAINTIES

The limits in table I mostly assume that the underlying cosmological model is the standard spatially flat Λ CDM model with adiabatic primordial perturbations. Slight variants have been considered, e.g. running spectral index of the primordial perturbation spectrum [8] and varying equation of state parameter w for the dark energy component [6, 9]. A significant degeneracy between w and M_ν was pointed out by Hannestad [9]. This degeneracy is a result of the fact that the matter power spectrum depends on $f_\nu = \Omega_\nu/\Omega_m$, and allowing w to vary weakens the constraints on Ω_m and hence indirectly on $M_\nu \propto f_\nu \Omega_m$. The degeneracy can be broken by including constraints from e.g. baryon acoustic oscillations [10] as shown in [6].

The spatially flat Λ CDM model describes the existing cosmological data well, but we are not yet in a position to exclude significant variations. For example, models where gravity is different from standard Einstein gravity are still viable, and that might change the neutrino mass limit significantly (see e.g. [11]) for an example).

The CMB is a very clean cosmological probe in the sense that the extraction of the anisotropy signal from the data involves relatively few and well justified

*Electronic address: oelgaroy@astro.uio.no

†Electronic address: lahav@star.ucl.ac.uk




FIG. 1: The power spectrum of the full 2dFGRS (bars) along with models with $f_\nu = 0$, $\Omega_m h = 0.168$ (full line) and $f_\nu = 0.18$, $\Omega_m h = 0.38$ (dashed line).

astrophysical assumptions. From this point of view the upper mass bound of 2 eV at the 95 % confidence level from the CMB data alone found by Fukugita et al. [7] is the most robust one. However, if one extends the space of models investigated to models that are very different from Λ CDM, there is no upper bound on neutrino masses from the CMB. To demonstrate this point, we need just to point out that Blanchard et al. [12] found that an Einstein-de Sitter model with $M_\nu = 2.4$ eV gave an excellent fit to the WMAP data, provided that there are oscillations in the primordial power spectrum, as produced e.g. if there is a phase transition during inflation. Now this model has both a low Hubble parameter ($h = 0.46$), no cosmological constant (and hence a bad fit to the supernovae type Ia data), and is also in some tension with the baryonic acoustic oscillations, but this goes to show that the CMB alone cannot produce a constraint on the neutrino mass once one allows for more radical departures from Λ CDM.

To get really tight constraints on the neutrino masses, one needs to include large-scale structure data, since the

CMB alone cannot go much below 2 eV in sensitivity. It is, however, worth nothing that large-scale structure alone cannot do the job. Take the 2dFGRS power spectrum as an example. Figure 1 below shows the power spectrum of the full 2dFGRS survey as determined by Cole et al [13]. Figure 1 also shows the power spectra for a model with no massive neutrinos and $\Omega_m h = 0.168$, and for a model with $f_\nu = 0.18$ and $\Omega_m h = 0.38$. These two models have identical values of the χ^2 for the data in the range used in fits, marked by the dashed vertical lines in the figure. Thus, one can still hide a lot of neutrinos in the matter power spectrum when one does not make use of external constraints on Ω_m .

IV. ASTROPHYSICAL SYSTEMATICS

Galaxy redshift surveys measure the distribution of galaxies in the local universe. The relation between the distribution of the luminous matter and the dark matter is therefore an important issue when estimating cosmological parameters in general, and the neutrino mass in particular (see e.g. [14] for an overview). It is well known that red galaxies are more common in the centres of rich clusters than blue galaxies, and one can get a simple, empirical estimate of the importance of biasing by doing the neutrino mass analysis separately for red and blue galaxies. An investigation of this effect is in preparation [15], and the preliminary results indicate that the effect is potentially important.

V. SUMMARY

Current cosmological observations provide strong upper limits on the sum of the neutrino masses. However, when assessing the significance of these limits one should bear in mind that several assumptions are involved in deriving these limits, both cosmological and astrophysical. It is an important task for further research to clarify how sensitive the results are to these assumptions, and how well they are justified.

-
- [1] Spergel, D. N. *et al.*, arXiv:astro-ph/0603449
 - [2] Mangano, C., Miele, G., Pastor, S., Pinto, T., Pisanti, O. and Serpico, P. D., Nucl. Phys. B **729**, 221 (2005).
 - [3] Lesgourgues, J. and Pastor, S., arXiv: astro-ph/0603494
 - [4] Elgarøy, Ø. *et al.*, Phys. Rev. Lett. **89**, 061301 (2002).
 - [5] Sanchez A., Baugh C. M., Percival W. J., Peacock J. A., Padilla, N. D., Cole, S., Frenk, C. S. and Nordberg, P., Mon. Not. Roy. Astron. Soc. **366**, 189 (2006).
 - [6] Goobar, A., Hannestad, S., Mortsell, E. and Tu, H., arXiv:astro-ph/0602155.
 - [7] Fukugita, M., Ichikawa, K., Kawasaki, M. and Lahav, O., arXiv: astro-ph/0605362.
 - [8] Seljak, U., Slosar, A. and McDonald, P., arXiv:astro-ph/0604335.
 - [9] Hannestad, S., Phys. Rev. Lett. **95**, 221301 (2005).
 - [10] Eisenstein, D. *et al.*, Astrophys. J. **633**, 560 (2005).
 - [11] Skordis, C., Mota, D. F., Ferreira, P. G. and Boehm, C., Phys. Rev. Lett. **96**, 011301 (2006).
 - [12] Blanchard, A., Douspis, M., Rowan-Robinson, M. and Sarkar, S., Astron.Astrophys. **412**, 35 (2003).
 - [13] Cole, S. *et al.*, Mon. Not. Roy. Astron. Soc. **362**, 505 (2005).
 - [14] Elgarøy, Ø. and Lahav, O., JCAP **0304**, 004 (2003).
 - [15] Elgarøy, Ø., Lahav, O., Percival, W. J., Cole, S. and Peacock, J. A., in preparation.